

## SAFETY PRECAUTIONS

### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when applying via hand-held equipment. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

### Environmental protection:

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from broadcast air-assisted sprayers to fall within 20m of the top of the bank of any static or flowing waterbody when applied to apple, pear and cherry; within 10m when applied to blackcurrant, redcurrant, gooseberry and blackberry; within 5m when applied to grape. Or within 5m of a ditch which is dry at the time of application of the top of the bank of a static or flowing waterbody, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5m of the top of a ditch which is dry at the time of application. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. For crops with 5m buffer zone only: DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water. ALL CROPS WITH A BUFFER ZONE GREATER THAN 5M ARE NOT ELIGIBLE FOR BUFFER ZONE REDUCTION UNDER THE LERAP HORIZONTAL BOOM SPRAYERS SCHEME.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer or a broadcast air-assisted sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone as appropriate to the crop must be maintained. The results of the LERAP must be recorded and kept available for three years.

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

# Landseer

L I M I T E D

## Systhane® 20 EW

### FUNGICIDE

TM ® Trademark of Corteva Agriscience and its affiliated companies.

Product Registration Number: MAPP 19160

GROUP 3 FUNGICIDE

An oil in water emulsion containing 200 g/litre (20% w/w) myclobutanil.

A systemic fungicide for the control of APPLE SCAB, PEAR SCAB and POWDERY MILDEW in APPLES and PEARS, BROWN ROT in CHERRY, AMERICAN POWDERY MILDEW in CURRANTS, GOOSEBERRIES, and BLACKBERRIES, POWDERY MILDEW in STRAWBERRIES and POWDERY MILDEW in GRAPES

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

**READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.**

**PROTECT FROM FROST.**



Sythane® 20EW; Myclobutanil

### WARNING

**Causes serious eye irritation.**

**Suspected of damaging the unborn child.**

**May cause damage to organs through prolonged or repeated exposure.**

**Toxic to aquatic life with long lasting effects.**

Do not handle until all safety precautions have been read and understood.

Use personal protective equipment as required

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Get medical advice/attention if you feel unwell.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste

**To avoid risks to human health and the environment, comply with the instructions for use.**

**Authorisation Holder**  
Corteva Agriscience UK Limited

CPC2 Capital Park, Fulbourn,  
Cambridge, CB21 5XE.

Telephone: (01462) 457272

24 Hour Emergency Telephone Number:

0161 884 1235

National Poisons Information Service : 111

Technical Hotline: 0800 689 8899 or

email [ukhotline@corteva.com](mailto:ukhotline@corteva.com)

**Distributed by:**

**Landseer Limited**

Lodge Farm, Goat Hall Lane,  
Galleywood, Chelmsford,

Essex CM2 8PH

Tel: (01245) 357109

Fax: (01245) 494165



The Voluntary Initiative

## NET CONTENTS : 1 LITRE e

This label is compliant with the CPA Voluntary Initiative Guidance.



IMPORTANT INFORMATION					
FOR USE ONLY AS A PROFESSIONAL FUNGICIDE					
Crop/situations	Maximum Individual Dose	Maximum Total Dose	Maximum number of treatments	Latest Time of Application	Aquatic buffer zone distance (metres):
Apple, pear	0.45 litres product/ha (see Other Specific Restrictions)	1.35 litres product/ha/annum	3	14 days before harvest	20 m
Cherry	0.45 litres product/ha	0.9 litres product/ha/annum	2	7 days before harvest	20 m
Blackcurrant and redcurrant (outdoor), gooseberry (outdoor), blackberries (outdoor)	0.45 litres product/ha	1.0 litre product per hectare per crop	3	14 days before harvest	10 m
Strawberry (outdoor)	0.5 litres product/ha	1.0 litres product/ha/annum	2	14 days before harvest	-
Strawberry (protected)	0.3 litres product/ha	0.9 litres product/ha/annum	3	3 days before harvest	-
Grape (wine)	0.24 litres product/ha	0.72 litres product/ha/annum	3	14 days before harvest	5 m
<b>Other Specific Restrictions:</b> Only 1 application may be made to apple, pear and cherry crops during August. DO NOT HANDLE TREATED GRAPEVINES for at least 12 days after treatment.					
<b>READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.</b>					

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

SYSTHANE 20EW with the active Myclobutanil is a systemic fungicide of the triazole class providing protectant activity. Systhane 20EW must be used as a preventative application, alone or in combination with a contact fungicide or as part of a seasonal spray program. Myclobutanil is a triazole fungicide, which acts by inhibiting sterol biosynthesis in fungi.

## TAINT TESTING

Taint tests have shown that SYSTHANE 20EW does not taint apple and currants, but growers should consult processors before use on any other crop.

## RESISTANCE

SYSTHANE 20EW contains a DMI fungicide. Sensitivity shifts to some DMI fungicides has been identified in Apple Scab (*Venturia inaequalis*) and powdery mildew (*Podosphaera leucotricha*) in apple and pear which may affect the performance of some products.

As part of an effective resistance risk management strategy Systhane 20 EW must be applied as a preventative application at the full label rate to ensure full crop coverage. Use Systhane 20EW in mixture or alternate with a different mode of action product at a rate which is effective on the target pathogen.

For further advice on resistance management in DMIs contact your agronomist or specialist advisor, and visit the FRAG-UK website.

## IPM PROGRAMMES

The effect of Systhane 20EW on pathogenic fungi based methods of IPM control has not been investigated, therefore use of Systhane 20EW in these situations is done so at the growers discretion.

## MIXING

**SYSTHANE 20EW Alone:** Shake the container well before use. Partially fill spray tank with water and start agitation. Pour required amount of Systhane 20EW into the tank and fill to required level. Maintain agitation during mixing and loading and until spraying is complete.

Tank mixing with PP Captan 80 WG: Pour the SYSTHANE 20EW into the partially filled spray tank as described above. Remove the filter basket and pour the required amount of PP Captan 80 WG directly into spray tank. DO NOT add the PP Captan 80 WG in a sudden large quantity. Agitate whilst topping up the tank and continue agitation before and during spraying.

## APPLES AND PEARS

Sythane 20EW is a systemic fungicide with protectant activity for the moderate control of apple scab (*Venturia inaequalis*), pear scab (*Venturia pirina*) and powdery mildew (*Podosphaera leucotricha*).

Sythane 20EW is most effective when used as part of a routine preventative spray programme from bud burst to the onset of petal fall. SYSTHANE 20EW is safe to use during flowering. From the onset of petal fall SYSTHANE 20EW should be tank-mixed with protectant fungicides to enhance fruit scab control. From mid June, SYSTHANE 20EW can be applied alone for the moderate control of secondary mildew.

SYSTHANE 20EW is safe to use on all the main commercial varieties of apples and pears grown in the UK. For PP Captan 80 WG mixtures with SYSTHANE 20EW please consult the PP Captan 80 WG label for details of varietal restrictions.

## SPRAY TIMING AND RATES OF USE

To help prevent leaf scab and reduce the development of powdery mildew from primary infections, apply SYSTHANE 20EW from bud burst to the onset of petal fall. To improve control of fruit scab after the onset of petal fall, tank mix SYSTHANE 20EW with PP Captan 80 WG (see COMPATIBILITY and MIXING).

Minimum spray interval	Bud burst to petal fall	From onset of petal fall
		Tank mix the listed rates (per ha) of Systhane 20EW with either
	Sythane 20EW	PP Captan 80WG
10 days	0.34 litres / ha	0.85 kg
10-14 days	0.45 litres / ha	1.2 kg

In periods of rapid growth or high disease pressure, SYSTHANE 20EW and partner fungicides should be used at 10 day instead of 10-14 day intervals.

Pest	Apple powdery mildew ( <i>Podosphaera leucotricha</i> ), apple scab ( <i>Venturia inaequalis</i> ) and pear scab ( <i>Venturia pirina</i> )
Level of control	Moderate control of powdery mildew ( <i>Podosphaera leucotricha</i> ). Moderate control of scab ( <i>Venturia inaequalis</i> and <i>Venturia pirina</i> ).
Rate	0.34 l/ha (10 day interval) 0.45 (10-14 day interval)
Water volume	200-500 litres per hectare
Maximum number of applications	3
Time of application	Apply SYSTHANE 20EW from the beginning of bud burst. Where disease pressure is high, use the shorter spray interval for optimum disease control.
Latest time of application	14 days before picking fruit

## STONE FRUIT (CHERRY)

Pest	Brown Rot ( <i>Monilia fructicola</i> and <i>laxa</i> )
Level of control	Moderate control of <i>Monilia fructicola</i> and <i>Monilia laxa</i>
Rate	0.45 l/ha
Water volume	500-1000 litres per hectare
Maximum number of applications	2
Time of application	SYSTHANE 20EW can be applied from bud burst. 10 days minimum should elapse before a further application of SYSTHANE 20EW is made.
Latest time of application	7 days before picking fruit

## CURRANTS (BLACK & RED), GOOSEBERRIES AND BLACKBERRIES

### CURRANTS (BLACK & RED), GOOSEBERRIES AND BLACKBERRIES

SYSTHANE 20EW is most effective when used as part of a routine preventative spray programme from just before the first signs of mildew infection up to two weeks before picking if necessary.

Pest	Powdery Mildew ( <i>Sphaerotheca mors-uvae</i> )
Level of control	Moderate control of powdery mildew ( <i>Sphaerotheca mors-uvae</i> ).
Rate	0.45 l/ha
Water volume	400-1000 litres per hectare
Maximum number of applications	3
Time of application	Apply SYSTHANE 20EW from the first leaves unfolded for currants and white bud stage for gooseberry. First application should be made at early infection at a minimum of 10 day intervals
Latest time of application	14 days before picking fruit

## STRAWBERRIES

SYSTHANE 20EW should be applied pre-harvest, beginning at or just prior to first flower and thereafter at the recommended spray intervals up to 3 days (protected) or 14 days (outdoor) before picking. Post-harvest sprays may be required where mildew is present and likely to be damaging (especially mildew susceptible varieties). This will help to produce vigorous growth and reduce the amount of over-wintering inoculum.

## STRAWBERRIES – PROTECTED USE

### SPRAY TIMING AND RATE OF USE

Pest	Powdery Mildew ( <i>Sphaerotheca macularis</i> )
Level of control	Moderate control of powdery mildew ( <i>Sphaerotheca macularis</i> )
Rate	0.3 l/ha
Water volume	400-1000 litres per hectare
Maximum number of applications	3
Time of application	Apply SYSTHANE 20EW from the first set of flowers at the bottom of the rosette. First application should be made at early infection at a minimum of 10 day intervals
Latest time of application	3 days before harvest

## STRAWBERRIES – OUTDOOR USE

### SPRAY TIMING AND RATE OF USE

Pest	Powdery Mildew ( <i>Sphaerotheca macularis</i> )
Level of control	Moderate control of powdery mildew ( <i>Sphaerotheca macularis</i> )
Rate	0.5 l/ha
Water volume	1000-2000 litres per hectare
Maximum number of applications	2
Time of application	Apply SYSTHANE 20EW from the first set of flowers at the bottom of the rosette. First application should be made at early infection at a minimum of 10 day intervals
Latest time of application	14 days before harvest

## GRAPE (WINE)

### SPRAY TIMING AND RATE OF USE

Pest	Powdery mildew ( <i>Uncinula necator</i> )
Rate	0.06 – 0.24 l/ha
Water volume	400-1600 litres per hectare
Maximum number of applications	3
Time of application	Apply SYSTHANE 20EW from the first flower hoods detached stage at a minimum 10 day interval between applications. Adjust rate to match growth stage and canopy development.
Latest time of application	14 days before harvest

## TANK CLEANING

All spraying equipment must be thoroughly cleaned both inside and out using a proprietary spray tank cleaner as follows:

1. Immediately after spraying thoroughly and completely rinse all internal surfaces.
2. Rinse inside of tank with clean water and flush through booms and in-line strainers using at least one tenth of the spray tank volume. Drain tank completely.
3. Fill tank with clean water (minimum 10% of tank capacity) and add tank cleaner at the recommended rate and agitate for 15 minutes. Flush the boom and hoses and drain tank completely.
4. Nozzles and filters should be removed and cleaned separately with a tank cleaner at the recommended rate.
5. Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume. Drain tank completely.
6. For disposal of washings, follow Code of Practice for Using Plant Protection Products. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4. Follow specific tank cleaning advice required for any partner products used in mixture with Systhane 20EW.

## COMPATIBILITY

SYSTHANE 20EW is compatible with PP Captan 80 WG. For compatibility information on other products please contact your dealer or advisor.

## COMPANY ADVISORY INFORMATION

The following information does not form part of the product label under the UK plant protection products regulations and provides additional advice on product use at the discretion of Corteva Agriscience.

## TRADEMARK ACKNOWLEDGEMENTS

SYSTHANE 20 EW is a trademark of Corteva Agriscience and its affiliated companies. All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

## CORTEVA AGRISCIENCE CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

For further information please contact Corteva Agriscience.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<https://www.corteva.co.uk/content/dam/dpagco/corteva/eu/gb/en/files/sds/SYSTHANE-SDS.pdf>

Alternatively contact your supplier

THIS PAGE HAS BEEN LEFT  
BLANK INTENTIONALLY

SPECIALS  
RECOMMEN  
THIS PAGE HAS BEEN LEFT  
BLANK INTENTIONALLY

#### SAFETY PRECAUTIONS

##### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. WEAR SUITABLE PROTECTIVE GLOVES when applying via hand-held equipment. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection. WASH CONCENTRATE from skin or eyes immediately. WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

##### Environmental protection:

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from broadcast air-assisted sprayers to fall within 20m of the top of the bank of any static or flowing waterbody when applied to apple, pear and cherry; within 10m when applied to blackcurrant, redcurrant, gooseberry and blackberry; within 5m when applied to grape. Or within 5m of a ditch which is dry at the time of application of the top of the bank of a static or flowing waterbody, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5m of the top of a ditch which is dry at the time of application. Aim spray away from water.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. For crops with 5m buffer zone only: DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water. ALL CROPS WITH A BUFFER ZONE GREATER THAN 5M ARE NOT ELIGIBLE FOR BUFFER ZONE REDUCTION UNDER THE LERAP HORIZONTAL BOOM SPRAYERS SCHEME.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer or a broadcast air-assisted sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone as appropriate to the crop must be maintained. The results of the LERAP must be recorded and kept available for three years.

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

##### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.  
KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.  
RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

# Landseer

LIMITED

## Systhane® 20 EW

### FUNGICIDE

™ Trademark of Corteva Agriscience and its affiliated companies.

Product Registration Number: MAPP 19160

GROUP 3 FUNGICIDE

An oil in water emulsion containing 200 g/litre (20% w/w) myclobutanil.


A systemic fungicide for the control of APPLE SCAB, PEAR SCAB and POWDERY MILDEW in APPLES and PEARS, BROWN ROT in CHERRY, AMERICAN POWDERY MILDEW in CURRANTS, GOOSEBERRIES, and BLACKBERRIES, POWDERY MILDEW in STRAWBERRIES and POWDERY MILDEW in GRAPES

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.

NET CONTENTS : 1 LITRE e



**Systhane® 20EW; Myclobutanil**  
**WARNING**  
Causes serious eye irritation.  
Suspected of damaging the unborn child.  
May cause damage to organs through prolonged or repeated exposure.  
Toxic to aquatic life with long lasting effects.  
Do not handle until all safety precautions have been read and understood.  
Use personal protective equipment as required  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
Get medical advice/attention if you feel unwell.  
Dispose of contents/container to a licensed waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste  
To avoid risks to human health and the environment, comply with the instructions for use.

**Authorisation Holder**  
Corteva Agriscience UK Limited  
CPC2 Capital Park, Fulbourn,  
Cambridge, CB21 5XE.  
Telephone: (01462) 457272  
24 Hour Emergency Telephone Number:  
0161 884 1235  
National Poisons Information Service : 111  
Technical Hotline: 0800 689 8899 or  
email ukhotline@corteva.com

**Distributed by:**  
Landseer Limited  
Lodge Farm, Goat Hall Lane,  
Galleywood, Chelmsford,  
Essex CM2 8PH  
Tel: (01245) 357109  
Fax: (01245) 494165



This label is compliant with the CPA Voluntary Initiative Guidance.