

SAFETY PRECAUTIONS

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), RUBBER BOOTS, SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.
WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces or applying the product.
WASH CONCENTRATE from skin or eyes immediately
DO NOT BREATHE SPRAY
WHEN USING DO NOT EAT, DRINK OR SMOKE
WASH HANDS AND EXPOSED SKIN before meals and after work
IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).
To protect aquatic organisms respect an unsprayed buffer zone of 4m to surface water bodies.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS
KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place
KEEP OUT OF REACH OF CHILDREN
EMPTY CONTAINER
COMPLETELY and dispose of safely
DO NOT RE-USE
CONTAINER for any purpose.



A 6 0008844



Zorvec Entecta™

FUNGICIDE

PCS 06766

Zorvec Entecta™ is a suspo emulsion formulation containing 240 g/L amisulbrom and 48 g/L oxathiapiprolin for the control of late blight in potatoes.

GROUP	21	49	FUNGICIDES
-------	----	----	------------

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.
PROTECT FROM FROST.

Authorisation Holder
Corteva Agriscience UK Limited
CPC2 Capital Park, Fulbourn, Cambridge,
Cambridgeshire. CB21 5XE.
Telephone: (01462) 457272
For Technical Enquires Call: +44 (0) 800 6898899 or email:ukhotline@corveva.com
24 Hour Emergency Telephone Number: 076 680 5288
National Poison Centre (Dublin): 01 8092166

™ Trademarks of Corteva Agriscience and its affiliated companies

Safety Information

ZORVEC ENTECTA

Contains 240 g/L amisulbrom and 48 g/L oxathiapiprolin

UFI: 19UA-KONQ-Y006-Q8NV

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/protective clothing/eye protection/face protection
IF exposed or concerned: Get medical advice / attention.
Store locked up
Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

Contains reaction mass of 5-chloro-2-methyl-2H-isothiazol-3-one and 2-methyl-2H-isothiazol-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS: 06766



WARNING



IMPORTANT INFORMATION

FOR USE ONLY AS AN PROFESSIONAL FUNGICIDE

Crop	Max. Individual Dose	Max. total dose	Max. No. of applications	Latest time of Application
Potato	0.25 L	1l.ha	4 per crop	7 days pre-harvest*

* See Directions for Use

Method of application: Tractor mounted / trailed sprayer

Other Specific Restrictions: Apply from BBCH 21-89

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE



To access the Safety Data Sheet for this product scan the QR code or use the weblink below:
<https://www.corteva.ie/content/dam/dpago/corteva/ie/en/files/sds/ZORVEC-ENTECTA-IRL-SDS.pdf>



3 576960 073899 >

NET CONTENTS: 1 LITRE e

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

Use in sequence of no more than 3 consecutive applications, or in alternation with a different mode of action fungicide.

Apply preventively before first visible signs of disease, using a 7 to 10-day spray interval. Use short interval in case of high disease pressure.

A minimum interval of 7 days between applications must be observed.

DISEASE CONTROL

Zorvec Entecta™ should be used as part of a season long programme of correctly timed sprays commencing before disease infection. When used as a preventative treatment, Zorvec Entecta™ gives control of late blight caused by *Phytophthora infestans*. Products from the same chemical group with similar mode of action to oxathiapiprolin must not be applied for more than 1/3 of total applications in the season, or 3 to 4 applications depending on the crop, whichever is the more restrictive.

According to Good Agricultural Practice, Zorvec Entecta™ should be applied on dry foliage. Use sufficient water to obtain thorough, uniform crop coverage.

Apply preventatively before the first visible signs of disease in the crop. Commence treatment with ZORVEC ENTECTA at the first blight warning or when local weather conditions are favourable for disease development. Applications should be repeated at 7-10 day spray intervals depending on disease pressure. As disease pressure increases, the interval should be shortened.

RESISTANCE

Repeated use of products for control of specific plant pathogens may lead to the selection of resistant strains of fungi and result in a reduction of disease control. A disease management program that includes rotation and/or applying mixtures of fungicides with different effective modes of action is essential to reduce the risk of fungicide resistance development.

The mode of action of oxathiapiprolin, one of the active ingredients in Zorvec Entecta™, is novel and has been assigned the FRAC code 49. Oxathiapiprolin inhibits an oxysterol binding protein (OSBP) homologue.

Amisulbrom, the other active ingredient, is a Qil-fungicide (Quinone inside Inhibitors). This group is considered to be medium to high risk for resistance.

The following restrictions apply to the use of Zorvec Entecta™ or any other FRAC 49-containing fungicides:

Exposure to Zorvec Entecta™ should not exceed thirty-three percent (33%) of the total period of protection needed per crop.

For potatoes when three (3) or more fungicide applications are made, use Zorvec Entecta™ in no more than 33% of the total number of fungicide applications with a maximum of four (4) applications per crop. Where less than three (3) fungicide applications are made, make no more than one (1) application of Zorvec Entecta™.

Zorvec Entecta™ applications are to be made preventively and no more than three (3) times in sequence before applying a fungicide with a different mode of action.

Where a fungicide with a different mode of action follows Zorvec Entecta™ application(s), this fungicide preferably should have curative activity.

There will be no more than six applications of oxathiapiprolin per year on the same acreage, targeting the same pathogen.

For guidance on a particular crop and disease control situation, consult your agronomist, crop protection specialist or Corteva Agriscience.

MIXING

Spraying equipment should be thoroughly cleaned before spraying. ZORVEC ENTECTA mixes easily with water and should be added directly to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is complete.

WATER VOLUME AND APPLICATION

Apply ZORVEC ENTECTA in at least 200 L water/hectare using a MEDIUM quality spray (as defined by BCPC) to give good even coverage of the crop. Higher water volumes should be used in dense crops to improve coverage.

COMPATIBILITY

For details on compatibilities contact your distributor or Corteva technical hotline on +44 (0) 800 689 8899 or ukhotline@corteva.com.

Zorvec Entecta™ is compatible with a range of fungicides and insecticides.

SPRAY TANK CLEANING

Immediately after application, completely drain spray equipment. Thoroughly rinse sprayer and flush the hoses, boom nozzles with clean water.

Loosen and physically remove visible deposits.

Remove and clean nozzles, screens and strainers.

Flush the entire system with clean water.

Take all necessary safety precautions when cleaning equipment.

Do not clean near wells or water sources.

Consult label tank cleanup procedures for all tank-mix partners and be sure to use the most rigorous procedure recommended.

Dispose of waste rinse water in accordance with local regulations.

TRADEMARK ACKNOWLEDGEMENTS

Zorvec Entecta is a trademark of Corteva Agriscience and its affiliated companies. All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

COMPANY ADVISORY INFORMATION

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SPECIMEN

THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY



Zorvec Entecta™

FUNGICIDE

PCS 06766

Zorvec Entecta™ is a suspo emulsion formulation containing 240 g/L amisulbrom and 48 g/L oxathiapiprolin for the control of late blight in potatoes.

GROUP	21	49	FUNGICIDES
-------	----	----	------------

**READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.
PROTECT FROM FROST.**

Authorisation Holder
Corteva Agriscience UK Limited
CPC2 Capital Park, Fulbourn, Cambridge,
Cambridgeshire. CB21 5XE.
Telephone: (01462) 457272
For Technical Enquires Call: +44 (0) 800 6898899 or email: ukhotline@corteva.com
24 Hour Emergency Telephone Number: 076 680 5288
National Poison Centre (Dublin): 01 8092166

™® Trademarks of Corteva Agriscience and its affiliated companies

NET CONTENTS: 1 LITRE

SAFETY PRECAUTIONS
Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), RUBBER BOOTS, SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.
WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces or applying the product.
WASH CONCENTRATE from skin or eyes immediately
DO NOT BREATHE SPRAY
WHEN USING DO NOT EAT, DRINK OR SMOKE
WASH HANDS AND EXPOSED SKIN before meals and after work
IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection
Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).
To protect aquatic organisms respect an unsprayed buffer zone of 4m to surface water bodies.

Storage and disposal
KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS
KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place
KEEP OUT OF REACH OF CHILDREN
EMPTY CONTAINER COMPLETELY and dispose of safely
DO NOT RE-USE CONTAINER for any purpose.



Safety Information
ZORVEC ENTECTA
Contains 240 g/L amisulbrom and 48 g/L oxathiapiprolin
UFI: 19UA-KONQ-Y006-Q8NV

Suspected of causing cancer.
Very toxic to aquatic life with long lasting effects.
Wear protective gloves/protective clothing/eye protection/face protection
IF exposed or concerned: Get medical advice / attention.
Store locked up
Collect spillage.
Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

Contains reaction mass of 5-chloro-2-methyl-2H-isothiazol-3-one and 2-methyl-2H-isothiazol-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS: 06766

WARNING

IMPORTANT INFORMATION
FOR USE ONLY AS AN PROFESSIONAL FUNGICIDE

Crop	Max. Individual Dose	Max. total dose	Max. No. of applications	Latest time of Application
Potato	0.25 L	1l.ha	4 per crop	7 days pre-harvest*

* See Directions for Use

Method of application: Tractor mounted / trailed sprayer
Other Specific Restrictions: Apply from BBCH 21-89

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:
<https://www.corteva.ie/content/dam/dpago/corteva/eu/ie/en/files/sds/ZORVEC-ENTECTA-IRL-SDS.pdf>

