

Technical Information Sheet

Forefront T is for control of annual and perennial broad-leaved weeds in grassland for grazing only.

Key facts

Product Registration Number:	PCS 05395
Active Ingredient:	30 g/litre aminopyralid + 240 g/litre triclopyr
Pack size:	2.0 L
Formulation:	Oil in water emulsion
Maximum Individual Dose:	2.0 L/ha
Maximum Total Dose:	2.0 L/ha
Timing:	Only use between 1st March and 31st July. For applications made between 1st March and 31st May, 1 application may be made per calendar year. For applications made between 1st June and 31st July, only 1 application may be made in a 2-year period
Water Volumes:	200-300 L/ha
Spray Quality:	Medium as defined by BCPC

Key benefits

- Best available long-term control of broadleaved dock, curled dock, creeping thistle, spear thistle, common nettle, creeping buttercup, dandelion, ragwort and chickweed in established grassland.
- Forefront T does not require an anti-foaming agent.
- Grazing by sheep or cattle can resume 7 days after Forefront T application.
- A single well-timed application of Forefront T can control weeds for up to 18 months.
- Rainfast in 1 hour

Best use advice

- Forefront T should only be used on established grassland fields grazed by cattle or sheep.
- Grazing by cattle or sheep can resume 7 days after treatment or when the foliage of any poisonous weeds has died and become unpalatable.
- Applications to silage, hay and haylage fields should only be made after the final cut has been taken.
 - Treated grass must not be cut for silage, hay or haylage in the calendar year of treatment.
 - Silage, hay and haylage produced from the treated area in the following calendar year must stay on farm and manure from animals fed on this must stay on farm and be spread only onto agricultural grassland.
- For optimum control apply 2-3 weeks after cutting provided there is sufficient regrowth.
- Single application of 2.0 L/ha is recommended.
- The label prohibits the use of Forefront T via hand-held equipment.
- DO NOT mix with phenoxy herbicides – this may compromise translocation of Forefront T to the roots.
- Clover can be stitched in 4 months after application. Grass seed can be drilled 1 month after application.
- **Forefront T contains aminopyralid which carries restrictions on use of manure and following crops.**
 - Manure produced from animals fed from grass treated with Forefront T **must stay** on farm and should only be spread onto agricultural grassland.
 - Potatoes, sugar beet, fodder beet, vegetables, beans and other legumes can be planted in the second calendar year following Forefront T treatment.
- For optimum control, Forefront T should be applied to actively growing weeds at the appropriate growth stage:

Weed Controlled	Optimum timing at 2.0 L/ha	Weed Controlled	Optimum timing at 2.0 L/ha
Broadleaved dock	Rosette stage up to 250 mm high or wide	Common nettle	Up to 300 mm high
Curled dock	Rosette stage up to 250 mm high or wide	Creeping buttercup	Before flowering
Creeping thistle	Rosette stage up to 250 mm high	Dandelion	Before flower bud visible
Spear thistle	Rosette stage up to 250 mm high	Ragwort	Rosette stage up to 200 mm high

Visit us at corteva.ie

USE PLANT PROTECTION PRODUCTS SAFELY. Always read the label and product information before use.

For further information including warning phrases and symbols refer to label.

Triple rinse containers, puncture and invert to dry at time of use.

Corteva Agriscience, CPC2 Capital Park, Fulbourn, Cambridge CB21 5XE. Tel: +44 1462 457272.

Marketing Company: Whelehan Crop Protection, Suite 11/12 Bunkila Plaza, Bracetown Business Park, Clonee, Dublin 15 Ireland

Tel 00 353 1 5742410 Email cropprotection@tpwhelehan.ie

Forefront® T contains aminopyralid and triclopyr.

January 2023 – this version supersedes all previous versions

Weed Spectrum

Annual Weeds	Doxstar Pro 2L/ha	Envy 1.5L/ha	Forefront 1.2L/ha	Grazon Pro 60ml/10L	Pastor Trio 1L/ha	Thistlex 1L/ha
Bindweed (black)	◆ 6TL/50mm	◆ 4TL	◆	◆	◆ 4TL	◆
Bristly ox-tongue	◆		◆	◆	◆	
Charlock	◆ 2TL	200mm			4TL	
Chickweed	◆ B4	B4	◆	◆	100mm/B4	◆ <100mm
Cleavers	◆ B4/100mm	B4	◆		200mm	
Corn chamomile	◆	◆ 150mm	◆	◆	◆ 150mm	◆
Corn marigold		◆ 6TL		◆	◆	◆
Cranesbill	◆		◆	◆		
Dead-nettles	◆ 4TL/50mm	◆ 2TL	◆	◆	◆ 2TL	
Fat-hen	◆ 2TL	◆ 2TL	◆ 2TL	◆ 2TL	◆ 2TL	◆ 2TL
Fool's parsley	◆		◆	◆		
Forget-me-not	◆ 4TL	4TL	◆		4TL	
Fumitory	◆ 2TL/50mm	◆ 2TL	◆		◆ 2TL	
Groundsel		◆ 2TL	◆	◆	◆ 2TL	◆
Hemp-nettle	◆	◆ 6TL/100mm			◆ 4TL	
Himalayan balsam				◆		
Knotgrass	◆ 2TL	6TL	◆ 4TL	◆ 4TL	◆ 4TL	
Mayweeds	◆	<200mm	◆	◆	<200mm	◆ <4TL
Nettle (small)	◆ 100mm		◆			
Nightshade (black)	◆ 6TL/100mm	◆ <4TL	◆	◆	◆	
Orache			◆			
Pale persicaria	◆	◆ 2TL	◆	◆	◆ 2TL	
Poppy		<4TL	◆		◆ 4TL	
Redshank		◆ 2TL			◆ 2TL	
Scarlet pimpernel	◆		◆	◆		
Shepherd's-purse	◆ 2TL/50mm	<4TL	◆	◆	<4TL	
Speedwells			◆			
Spurrey		◆ 2TL			◆ 2TL	
Wild radish	◆ 2TL	◆ <6TL/80mm			◆ <4TL/50mm	
Yellow rattle			◆	◆		

Key			
◆	Good control	◆	Anecdotal or limited information
◆	Moderate control	T L	True leaf
◆	Some control	B4	Before flowering
◆	No control	☐	Seedling
◆	No Information		

Where we have knowledge of how our products might affect weeds we have detailed it in these tables. ◆ indicates information based on anecdotal or limited data, and is only indicative and should not be considered as a recommendation for use on the part of Corteva Agriscience. The user assumes full responsibility for use on these weeds.

Perennial Weeds	Doxstar Pro 2L/ha	Envy 1.5L/ha	Envy 2L/ha	Forefront 1.2L/ha	Grazon Pro 60ml/10L	Pastor Trio 1L/ha	Pastor Trio 2L/ha	Thistlex 1L/ha
Bindweed (field)	◆		◆	◆	◆		◆	
Bracken				◆ 1m				
Bramble	◆			◆				◆
Broom	◆			◆				
Burdock	◆			◆				
Buttercups	◆			B4	◆	From seed	Rosette	
Cinquefoil (creeping)				◆			◆	
Clover, trefoil	◆	◆	◆	◆	◆		◆	◆
Coltsfoot	◆			◆	◆		◆	
Cow parsley				◆	◆			
Daisy (common)	◆			◆ <2TL/25mm	◆			
Daisy (ox-eye)				◆			◆	
Dandelion	◆		B4	B4	◆		Rosette	◆
Docks	200mm	◆	◆ 200mm	Rosette <250mm		◆	200mm	◆
Gorse	◆							
Ground elder	◆			◆	◆			
Ground ivy	◆							
Hawthorn	◆			◆	◆			
Hemlock			◆	◆	◆		◆	
Hogweed (giant)	◆			◆	◆			
Horsetail (Equisetum)			◆		◆		◆	
Japanese knotweed				◆ <1m high	◆ <1m high			
Knapweed (common)			◆	◆	◆		◆	◆
Lesser celandine	◆			◆	◆			
Mallow (common)			◆	◆	◆		◆	
Medick (black)	◆			◆	◆			
Mugwort (common)	◆			◆	◆			
Nettle (common)	◆		◆	<300mm	B4		◆	◆
Old man's beard				◆				
Plantain (greater)	◆		◆	◆	◆		Rosette	
Plantain (ribwort)	◆		◆	◆	◆		Rosette	
Ragwort				Rosette <200mm	◆			
Rosebay willowherb	◆			◆	◆			
Rushes								
Self-heal				◆	◆			
Silverweed				◆	◆			
Sorrel (common)	◆		◆	◆	◆		◆	
Thistles				Rosette <250mm	Rosette 4-10TL 150mm	1TL	◆	Rosette 150-250mm
Vetch, tare	◆	◆	◆	◆	◆	◆	◆	◆
Yarrow	◆	◆	◆	◆	◆	◆	◆	
Yellow/Flag Iris				◆	◆			