

Technical Information Sheet

IRELAND

Peacog[™] is a new fungicide for the control of *Septoria tritici*, brown rust and yellow rust in Winter Wheat, Winter Triticale, Winter Spelt and Durum Wheat.

Peacog is broad-spectrum fungicide containing Fenpicoxamid, more commonly known as Inatrea Active. Peacog is part of a new class of chemistry called picolinamides which has a new site of action against fungi in cereals. Peacog must always be applied in mixture.

Key facts

FUNGICIDE

Product Registration Number:	Peacoq - PCS 06752 UFI code: 9059-M0XQ-2002-81QX		
Active Ingredient:	Peacoq - 50g/L Fenpicoxamid (Inatreq) (FRAC Code 21)		
Pack Size:	5L		
Formulation:	Peacoq - Emulsifiable Concentrate		
Maximum Single Dose Rate:	Peacoq - 2L/ha		
Maximum Total Dose per crop:	Peacoq - 2L/ha		
Maximum No. of Applications:	1 per crop per season		
Application Timing:	Beginning of stem elongation (GS 30)- end of flowering (GS 69)		
Water Volume:	100-300L/ha. But ideally use a minimum spray volume of 200l/ha in line with Corteva's 2023 Inatreq application advice. Spray pressure at 2-3 bar is recommended		
Aquatic buffer zone	Buffer zone is determined by the STRIPE Water Tool depending on rate and nozzle type		


Best Use Advice

- Peacog may be used on all commercial varieties of winter crops of wheat, triticale, durum wheat and spelt wheat.
- Peacog must always be used in mixture with a product with a different mode of action at a dose that offers robust control.
- Ideally apply at T2 to see maximum yield benefits but Peacog could alternatively be applied at T1 if preferred.
- Apply at the recommended rate of 2L/ha.
- Peacog will give the best results before disease has become established in the crop but offers outstanding control of Septoria both with persistent protectant and curative activity.
- Peacog offers broad spectrum, superior disease control which leads to longer green leaf retention and improved yield.
- Rainfastness is 1 hour for Peacoa.

Resistance Management

- Peacog MUST always be used in a mixture with a partner product which has a different mode of action.
- You must not apply more than one application of Peacog to any cereal crop in the same year.
- Peacoq should be used in accordance with the instructions for use for the target disease at the specific growth rate indicated.

Visit us at corteva.ie Technical Hotline: +44 (0) 800 689 8899 or email:ukhotline@corteva.com USE PLANT PROTECTION PRODUCTS SAFELY. Always read the label and product information before use.





Tank Mixing and Application

- Peacoq has been tested in many multi-way tank mixes with a variety of PGRs, herbicides, trace elements and other key fungicides.
- Products should only be tank mixed if each product can be applied with the label recommendations for its use.
- For a full tank mix list please visit the website- https://www.corteva.ie/tools-and-advice/tank-mixes.html
- Before applying Peacoq, please see our best practice application ad and resources to help optimise field operations.
 Visit www.corteva.ie/inatreqadvice or scan the QR code.

Disease Spectrum

Peacoq will give control to the following diseases in the following crops shown in the table below.

Disease	Winter Wheat	Winter Durum Wheat	Winter Triticale	Winter Spelt Wheat
Septoria Leaf Blotch	С	С	С	С
Yellow Rust	С	С	-	С
Brown Rust	С	С	-	С
				C=Control

STRIPE- Surface Water Tool for Reducing the Impact of Pesticides in the Environment

STRIPE is an initiative which allows farmers to reduce the size of mandatory untreated areas of land near water courses (buffer zones) while helping to protect aquatic life from pesticide contamination by reducing exposure.

Buffer zones are determined by the STRIPE water tool when ditches are present.

The buffer zone may be reduced by the use of low drift nozzles and application rate as

<u>Del The following rable</u>						
	Full Rate	¾ Rate	½ Rate			
Non- Drift reducing nozzles	30m	23m	15m			
50% Drift reducing nozzles	15m	11m	7m			
75% Drift reducing nozzles	7m	6m	4m			
90% Drift reducing nozzles	3m	2m	1m			

