

OcTtain™ XL

Trusted, easy to use, high performance broadleaf weed control.

HERBICIDE

WHY USE OCTTAIN™ XL HERBICIDE?

- Performance and value. The standard for kochia (including Group 2 and 9 resistant), plus wild buckwheat and 25 other key broadleaf weeds.
- Ease of use. Available in 40-acre cases, 240-acre drums and 1,280-acre totes.
- Crop-safe and flexible. Excellent crop safety in barley, wheat and durum.
- Tank mix with Simplicity™ GoDRI™ (wheat), Liquid Achieve™ (barley) herbicides and other grass weed control products.

BROADLEAF WEEDS CONTROLLED

- Annual sunflower
- Blue lettuce^{1,4}
- Bluebur
- Burdock
- Cleavers³
- Cocklebur
- Dandelion^{2,4}
- Docks⁴
- Dog mustard⁴
- Field bindweed^{1, 4}
- Field horsetail¹
- Field peppergrass⁴
- Flixweed

- Goat's-beard
- Gumweed⁴
- Hairy galinsoga⁴
- Hedge bindweed⁴
- Hemp nettle
- Hoary cress¹
- Kochia³
- Lamb's-quarters
- Leafy spurge^{1,4}
- Mustards (except green and grey tansy)
- Oak-leaved goosefoot⁴
- Plantain
- Prickly lettuce
- Ragweed

- Redroot pigweed⁴
- Round-leaved mallow
- Russian thistle⁴
- Shepherd's purse
- Smartweed⁴
- Stinkweed
- Stork's-bill
- Sweet clover
- Tansy mustard⁴
- Tartary buckwheat⁴
- Vetch
- Volunteer canola⁵
- Volunteer flax
- Wild buckwheat
- Wild radish

WEEDS SUPPRESSED

- Annual sow thistle⁴
- Canada thistle^{1,4}
- Chickweed³
- Perennial sow thistle¹
- Redroot pigweed

APPLICATION GUIDELINES

| | | |
|---|----------------------------|--|
| CROPS Barley Durum wheat Spring wheat Winter wheat | PACKAGING | Case: 2 x 9 L jugs Pallet: 5 x 108 L drums Tote: 576 L |
| | RATES | 450 ml/ac |
| | ACRES TREATED | • 20 ac/jug • 240 ac/drum • 1280 ac/tote |
| | WATER VOLUME | • Ground 20-40 L/ac (5-10 US gal/ac) • Aerial 12-20 L/ac (3-5 US gal/ac) |
| | TIMING | Crop stage: 4-leaf to just prior to flag leaf emergence Weed stage: 1 to 6-leaf |
| | RAINFAST | 1 hour |
| | TANK MIX HERBICIDES | <ul style="list-style-type: none"> • Assert® • Everest® • Horizon® • Liquid Achieve™ SC • Puma® Super • Simplicity™ • Simplicity™ GoDRI™ • Traxos® |



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Alfalfa
- Barley
- Canola
- Corn
- Dry beans
- Faba beans
- Flax
- Forage grasses
- Lentils
- Mustard
- Oats
- Peas
- Potatoes
- Rye
- Soybeans
- Sugar beets
- Sunflowers
- Wheat.

GRAZING AND HARVEST

- Allow 7 days after application before grazing lactating animals.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

¹ Top growth control only

² Spring rosettes

³ Including Group 2 resistant biotypes

⁴ Requires the addition of 86 ml/ac (2 oz/ac) of 2,4-D Ester 700

⁵ All herbicide-tolerant varieties