

Prominex™

Arylex™ active

HERBICIDE

Control annual AND perennial broadleaf weeds with the convenience of an all-in-one formulation.

WHY USE PROMINEX™ HERBICIDE IN WHEAT AND BARLEY?

- Delivers exceptional control of BOTH annual AND perennial broadleaf weeds in wheat and barley.
- With Arylex™ active you can **Just GO**. Small or large weeds, early or late crop stage and in cool or dry conditions.
- Flexibility to tank mix with the right graminicide and phenoxy partner.

PROMINEX ALONE

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Canada fleabane²
- Chickweed²
- Cleavers¹
- Common ragweed²
- Cow cockle
- Flixweed¹

- Giant ragweed²
- Hemp-nettle¹
- Henbit
- Kochia²
- Lamb's-quarters
- Nightshade species (*Eastern black, hairy and cutleaf*)
- Redroot pigweed
- Round-leaved mallow

- Shepherd's purse¹
- Stork's-bill
- Velvetleaf
- Volunteer alfalfa
- Volunteer flax
- Wild buckwheat
- Wild radish

GRASS WEEDS CONTROLLED

- Barnyard grass

PERENNIAL WEEDS CONTROLLED

- Canada thistle

WEEDS SUPPRESSED

- Annual sow thistle
- Wild mustard

PROMINEX + 235 ml/ac MCPA ESTER 600 (5 oz/ac)

ALL PROMINEX ALONE WEEDS PLUS:

- Annual sow thistle
- Ball mustard
- Burdock

- Cocklebur
- Field horsetail (*top growth*)
- False ragweed
- Plantain
- Prickly lettuce
- Stinkweed
- Vetch
- Volunteer canola
- Wild mustard
- Wild radish
- Wild sunflower

WEEDS SUPPRESSED

- Dandelion
- Smartweed
- Perennial sow thistle

PROMINEX + 241 ml/ac 2,4-D ESTER 700 (6 oz/ac)

ALL PROMINEX ALONE WEEDS PLUS:

- Annual sow thistle
- Bluebur

- Burdock
- Cocklebur
- False ragweed
- Goat's beard
- Plantain
- Prickly lettuce
- Russian thistle
- Stinkweed
- Volunteer canola
- Wild radish

- Wild sunflower

APPLICATION GUIDELINES

CROPS Barley Durum wheat Spring wheat Winter wheat	PACKAGING	Case: 2 x 8.3 L jugs Drum: 99.4 L
	RATE	415 ml/ac
	ACRES TREATED	• 20 ac/jug • 240 ac/drum
	TIMING	Crop stage: 3-leaf to just prior to flag leaf emergence
	RAINFAST	4 hours
	TANK MIX HERBICIDES	• Axial® • Everest® • Horizon • Liquid Achieve™ SC • Puma® Advance • Simplicity™ • Simplicity™ GoDRI™ • Traxos® • Trondus®



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Barley
- Canola
- Corn
- Flax
- Forage grasses
- Oats
- Mustard
- Peas¹
- Fall rye (*not underseeded with legumes, clover or alfalfa*)
- Soybeans¹
- Summerfallow
- Wheat

¹ Including Group 2 resistant biotypes
² Including Groups 2 & 9 resistant biotypes

¹ For pea or soybean rotation, rainfall from June 1 to August 31 in the year of application must be greater than 140 mm (5.5 inches) and annual rainfall must be greater than 175 mm (6.9 inches).