

Rezuvant™

Arylex™ active

HERBICIDE

Rezuvant™ herbicide combines leading Group 1 grass weed control with outstanding broadleaf technology in wheat and barley.

WHY USE REZUVANT™ HERBICIDE IN WHEAT AND BARLEY?

- Crop-safe, flexible and leading Group 1 performance on wild oats, green foxtail, barnyard grass and more.
- Unparalleled control of cleavers, hemp-nettle, wild buckwheat, kochia and many other tough broadleaf weeds.
- Tank mix with MCPA Ester 600 for expanded Group 4 mode of action control.

REZUVANT ALONE¹

BROADLEAF WEEDS CONTROLLED

- American dragonhead
- Canada fleabane
- Chickweed²
- Cleavers²
- Common ragweed
- Cow cockle
- Cutleaf nightshade
- Flixweed
- Hairy nightshade
- Hemp-nettle²

- Henbit
- Kochia
- Lamb's-quarters²
- Nightshade species (*Eastern black, hairy and cutleaf*)
- Redroot pigweed
- Round-leaved mallow²
- Shepherd's purse²
- Stork's-bill²
- Velvetleaf
- Volunteer alfalfa
- Volunteer flax²
- Wild buckwheat

GRASS WEEDS CONTROLLED

- Barnyard grass
- Green foxtail
- Proso millet
- Volunteer oats
- Volunteer canary seed
- Wild oats
- Yellow foxtail

WEEDS SUPPRESSED

- Annual sow thistle
- Wild mustard

REZUVANT + 235 ml/ac OF MCPA ESTER 600 (5 oz/ac)¹

ALL REZUVANT ALONE WEEDS PLUS:

- Annual sow thistle
- Ball mustard

- Burdock
- Cocklebur
- Plantain
- Prickly lettuce

- Ragweed (*common, false and giant*)
- Stinkweed
- Vetch
- Volunteer canola

- Wild mustard
- Wild radish
- Wild sunflower

APPLICATION GUIDELINES

CROPS Barley Spring wheat Winter wheat	PACKAGING Case: • Rezuvant A: 1 x 4.9 L jug • Rezuvant B: 2 x 10 L jugs Bulk: • Rezuvant A: 2 x 9.8 L jugs • Rezuvant B: 1 x 80 L drums
	RATES • Rezuvant A: 123 ml/ac • Rezuvant B: 500 ml/ac
	ACRES TREATED Case: 40 ac/case Bulk Unit: 160 acres
	WATER VOLUME Ground 20-40 L/ac (5-10 US gal/ac)
	TIMING Crop stage: 3-leaf to just prior to flag leaf emergence
	RAINFAST 1 hour
TANK MIX HERBICIDES MCPA Ester 600	

MIXING INSTRUCTIONS:

1. Fill sprayer 1/2 full of water and begin gentle sprayer tank agitation.
2. Add required amount of Rezuvant A and agitate for 2-3 minutes.
3. Add required amount of MCPA Ester 600 and agitate for 2-3 minutes.
4. Add required amount of Rezuvant B and agitate for 2-3 minutes.
5. Continue gentle agitation and add remainder of water.

CROP ROTATION

- Alfalfa
- Barley
- Canola
- Corn
- Dry bean (*Phaseolus vulgaris species including pinto, kidney and white types*)
- Faba beans
- Field peas
- Flax
- Mustard
- Oats
- Potatoes (*except seed potatoes*)
- Soybeans
- Spring wheat
- Sunflowers
- Timothy

GRAZING AND HARVEST

- Livestock may be grazed on treated crops 21 days following application.
- Do not harvest the treated crop within 60 days after application.

¹ For best weed control apply at these growth stages:

Grass - apply from 1-6 leaf, prior to emergence of the 4th tiller.

Broadleaf - apply from 1-8 leaf, unless otherwise specified (refer to product label).

² Overlapping Group 4 control of this weed when tank mixed with MCPA Ester at 5 oz per acre.