

Simplicity™

HERBICIDE

Superior performance, including elite Group 2 wild oat and bonus broadleaf weed control with no re-cropping restrictions.

WHY USE SIMPLICITY™ HERBICIDE?

- Superior performance. Elite grass and broadleaf weed control.
- Wide window of application. This is the only Group 2 wild oat product that can be applied up to just prior to flag leaf emergence.
- Resistance management. Controls Group 1 resistant wild oats.
- Tank mix flexibility.
- Rotational freedom to all major crops the following year.

BROADLEAF WEEDS CONTROLLED

- Chickweed¹
- Cleavers¹
- Corn spurry
- Cow cockle
- Flixweed
- Hemp-nettle¹
- Redroot pigweed
- Round-leaved mallow
- Shepherd's purse
- Smartweed¹
- Stinkweed
- Volunteer canola (excluding Clearfield[®])

GRASS WEEDS CONTROLLED

- Wild oats
- Barnyard grass
- Downy brome (fall application)
- Japanese brome
- Yellow foxtail

WEEDS SUPPRESSED

- Canada thistle
- Dandelion
- Downy brome (spring application)
- Green foxtail²
- Persian darnel
- Russian thistle
- White cockle
- Wild buckwheat

APPLICATION GUIDELINES

CROPS Durum wheat Spring wheat Winter wheat	PACKAGING • Case: • Simplicity: 2 x 8 L jugs • Water Conditioner: 2 x 1.5 L jugs
	RATES • Simplicity: 200 ml/ac • Water Conditioner: 37.5 ml/ac
	ACRES TREATED 40 ac/jug (80 ac/case)
	WATER VOLUME • Ground 12-40 L/ac (3-10 US gal/ac) • Aerial minimum 12 L/ac (3 US gal/ac)
	TIMING Crop stage: 3-leaf to just prior to flag leaf emergence Weed stage: • 1 to 4-leaf, two tillers stage of wild oats • 1 to 5-leaf stage of broadleaf weeds
	RAINFAST 2 hours
	TANK MIX HERBICIDES³ • Attain™ XC • Buctril® M • Cirpreme™ XC • Curtail® M • Exhilarate™ • MCPA Ester 600 • OcTtain™ XL • Prestige™ XL • Prominex™ • Pixxaro™ • Refine Extra® • Stellar™ XL • Thumper® • Tilt® fungicide



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Barley
- Brown mustard
- Canola
- Canola-quality brassica juncea
- Chickpea
- Dry bean (*Phaseolus vulgaris* species)
- Field peas
- Flax
- Lentils
- Oats
- Potatoes
- Soybean
- Spring wheat
- Sunflower
- Yellow mustard

GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Mature crops may be harvested 50 days after application.

¹ Not including group 2 resistant

² Corteva Agriscience research trials indicate that application to small stage, actively growing plants provides an increased level of control

³ Unless otherwise stated. Always read and follow label directions.