

Tandem™

HERBICIDE

Control problem grass and broadleaf weeds in wheat – with the tougher, easier, total-acre solution.

WHY USE TANDEM™ HERBICIDE?

- Tougher. Control of wild oats, Japanese brome, chickweed, cleavers, hemp-nettle, kochia, wild buckwheat and many more.
- Easy to use. Control all problem weeds using one solution.
- Flexible. Wide window of application, excellent crop safety, flexible rate and rotational freedom.
- Multi-effective mode of action. Two unique modes of action provide overlapping activity on hard-to-kill weeds for superior resistance management.

TANDEM ALONE

BROADLEAF WEEDS CONTROLLED

- Chickweed^{1,4}
- Cleavers^{1,4}
- Corn spurry
- Cow cockle
- Flixweed
- Hemp-nettle

- Kochia⁴
- Redroot pigweed
- Round-leaved mallow¹
- Shepherd's purse
- Smartweed
- Stinkweed
- Volunteer canola²
- Volunteer flax

GRASS WEEDS CONTROLLED

- Barnyard grass
- Japanese brome
- Wild oats²
- Yellow foxtail

WEEDS SUPPRESSED

- Canada thistle
- Dandelion⁵
- Downy brome⁴
- Green foxtail⁵
- Russian thistle
- Stork's-bill
- Wild buckwheat

TANDEM + 241 ml/ac OF 2,4-D ESTER 700 (6 oz/ac)

ALL TANDEM ALONE WEEDS PLUS:

- Bluebur
- Burdock
- Cocklebur

- Flixweed¹
- Goat's-beard
- Lamb's-quarters¹
- Plantain
- Prickly lettuce

- Ragweed
- Redroot pigweed¹
- Smartweed¹
- Stinkweed¹
- Sweet clover

- Volunteer sunflower
- Wild buckwheat
- Wild mustard¹
- Wild radish

TANDEM + 234 ml/ac OF MCPA ESTER 600 (5 oz/ac)

ALL TANDEM ALONE WEEDS PLUS:

- Burdock
- Cocklebur
- Lamb's-quarters¹

- Prickly lettuce
- Ragweed
- Shepherd's purse¹
- Smartweed¹

- Stinkweed¹
- Vetch
- Volunteer sunflower
- Wild buckwheat

- Wild mustard
- Wild radish

1 Weed controlled through multi-effective mode of action

2 Clearfield volunteer canola is controlled with the addition of MCPA or 2,4-D

3 Including Group 1 resistant biotypes

4 Including Group 2 resistant biotypes

5 Corteva Agriscience research trials indicate that application to small stage, actively growing plants provides an increased level of control

6 Spring rosettes

7 Top growth only

APPLICATION GUIDELINES

CROPS Durum wheat Spring wheat Winter wheat	PACKAGING	Case: • Tandem A: 1 x 8 L jug • Tandem B: 1 x 4.84 L jug • Water conditioner: 1 x 1.5 L jug
	RATES	• Tandem A: 200 ml/ac • Tandem B: 121 ml/ac • Water conditioner: 37.5 ml/ac
	ACRES TREATED	40 ac/case
	WATER VOLUME	Ground: 20-40 L/ac (5-10 US gal/ac) Aerial: Min. 12 L/ac (3 US gal/ac)
	TIMING	Crop stage: 3-leaf to just prior to flag leaf emergence Weed stage: 1 to 4-leaf, two tillers stage for wild oats and 1 to 5-leaf stage for broadleaf weeds
	RAINFAST	2 hours
	TANK MIX HERBICIDES	• Curtail® M herbicide • MCPA Ester 600 • 2,4-D Ester 700



DOWNLOAD THE 2023 FIELD GUIDE APP FOR PRODUCT SPECIFIC TANK MIX INSTRUCTIONS.

CROP ROTATION

- Barley
- Canola
- Chickpeas
- Corn
- Dry beans
- Flax
- Lentils
- Mustard
- Oats
- Peas
- Potatoes
- Soybeans
- Sunflower
- Wheat

GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application.
- Allow 7 days after application before grazing lactating animals.
- Withdraw meat animals from treated areas at least 3 days before slaughter.
- Allow 30 days after application before cutting hay or harvesting forage.
- Mature crops may be harvested 60 days after application.

Additional broadleaf weeds controlled when tank mixed with Curtail M herbicide (10 ac/jug)

- Annual sow thistle
- Burdock
- Canada thistle
- Cocklebur
- Common groundsel
- Dandelion⁴
- Field horsetail⁵
- Lamb's-quarters
- Perennial sow thistle
- Plantain⁵
- Prickly lettuce
- Ragweed
- Russian pigweed
- Scentless chamomile
- Sunflower (annual and volunteer)
- Tartary buckwheat
- Vetch
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild radish