

Tridem™

HERBICIDE

Delivers powerful broadleaf weed control combined with Group 2 grass chemistry for wheat farmers.

WHY USE TRIDEM™ HERBICIDE IN WHEAT?

- Delivers wide spectrum grass and broadleaf weed control for wheat farmers.
- Perennial broadleaf weed control without the re-cropping restrictions.
- With two different modes of action, Tridem provides overlapping control of certain key broadleaf weeds.

TRIDEM + 215 ml/ac OF 2,4-D ESTER 700 (5 oz/ac)

BROADLEAF WEEDS

- Annual sow thistle
- Burdock
- Bluebur
- Cleavers¹
- Cocklebur
- Common chickweed¹
- Cow cockle
- Daisy fleabane
- False flax
- Flixweed
- Goat's-beard

- Kochia
- Lamb's-quarters
- Narrow-leaved hawk's beard
- Plantain
- Prickly lettuce
- Ragweed
- Redroot pigweed¹
- Round-leaved mallow¹
- Russian pigweed
- Russian thistle
- Shepherd's purse¹
- Smartweed¹

- Stinging nettle
- Stinkweed¹
- Stork's-bill
- Sweet clover
- Thyme-leaved spurge
- Volunteer canola¹ (all herbicide tolerant biotypes)
- Volunteer flax¹
- Wild buckwheat¹
- Wild mustard¹
- Wild radish
- Wild sunflower¹

GRASS WEEDS CONTROLLED

- Japanese brome
- Wild oats

PERENNIAL WEEDS CONTROLLED

- Canada thistle
- Dandelion
- Perennial sow thistle

WEEDS SUPPRESSED

- Hemp-nettle

APPLICATION GUIDELINES

CROPS Durum wheat Spring wheat Winter wheat	PACKAGING Case: • Tridem A: 1 x 0.84 kg jug • Tridem B: 2 x 8.1 L jugs • Bindem: 1 x 2.4 L jug Bulk Pack: • Tridem A: 1 x 5.04 kg jug • Tridem B: 1 x 97.2 L drum • Bindem: 2 x 7.1 L jugs
	RATES • Tridem A: 21 g/ac • Tridem B: 405 ml/ac • Bindem: 60 ml/ac
	ACRES TREATED • 40 ac/case • 240 acre bulk pack
	TIMING Crop stage: 4-leaf to just prior to flag leaf emergence Weed stage: Apply early post-emergence to the main flush of actively growing weeds
	RAINFAST 2 hours
	TANK MIX HERBICIDES • 2,4-D Ester 700 • MCPA Ester 600

MIXING INSTRUCTIONS

1. Fill sprayer tank ½ full of water
2. Start sprayer tank agitation
3. Add the required amount of Tridem A Herbicide
4. Add the required amount of Tridem B Herbicide
5. Add 2,4-D Ester 700 next and continue agitation
6. Add required amount of Bindem utility modifier
7. Fill the sprayer tank with the remaining amount of water required

CROP ROTATION

- | | | | |
|-----------|---|---|-----------------------------------|
| • Alfalfa | • Dry common beans (phaseolus vulgaris) | • Mustard (brown, oriental and/or yellow) | • Potatoes (except seed potatoes) |
| • Barley | • Faba beans | • Oats | • Soybeans |
| • Canola | • Flax | • Peas | • Sunflower |
| • Corn | • Lentils | | • Wheat |

GRAZING AND HARVEST

Do not harvest grain within 60 days of application.

¹ Weed controlled by multi-effective mode of action when tank mixed with 2,4-D Ester at 5 oz/ac